



TITLE: Negative Pressure Wound Therapy for Patients with Diabetic Foot Ulcers and Pressure Ulcers: A Review of the Clinical Effectiveness

DATE: 21 June 2012

CONTEXT AND POLICY ISSUES

Diabetic foot ulcer (DFU) is a common diabetic complication.¹ From 2006 to 2008, it was reported that the prevalence for DFU in Medicare beneficiaries was about 8% in the United States.² DFU affects patients' quality of life significantly and also has a significant financial impact on healthcare system. A delay in management of DFU will increase morbidity and mortality and results in a higher amputation rate.³ Management of DFU is still a major therapeutic challenge throughout the world.⁴

Pressure ulcers (PU), also known as decubitus ulcers or bedsores, are caused by uninterrupted pressure on soft tissues overlying a bony prominence which obstructs the blood flow to the superficial tissues. The most commonly affected body parts are the sacrum or the hips, but other sites such as the elbows or knees can be affected.⁵ People with reduced mobility and poor skin condition in nursing homes or hospitals are more likely affected.⁵ Approximately 3 million adults in the United States suffer from pressure ulcers. The reported incidence of PU varies from 0.4% to 38.0%.⁶⁻⁸ The healing rates of the PU varies depending on the severity of PU, comorbidities and clinical management.⁶ The estimated costs of treatment for a PU ranges from \$37,800 to \$70,000. in the United States.^{6,7,9}

The management of DFU and PU is important to public health.^{1,4,10,11} Negative Pressure Wound Therapy (NPWT), also known as vacuum assisted closure therapy (VAC), topical negative pressure therapy or vacuum compression therapy^{1,12,13} appears to be an effective treatment for DFU and PU.^{10,14-17} The negative pressure can be delivered continuously or intermittently.¹⁸⁻²⁰ Most existing clinical evidence is derived from the use of open-cell polyurethane foam at -125 mmHg.²¹⁻²⁴ However, this pressure may cause pain and ischemia and often has to be reduced.^{23,24}

Recently, two new NPWT systems (Smith & Nephew PICO^{2,25} and KCI V.A.C.Via²⁶) have been released. These are less powerful NPWT systems that operate at lower pressures and are disposable after each use. More importantly, these newer machines are comparable in cost to standard wound care.

Disclaimer: The Rapid Response Service is an information service for those involved in planning and providing health care in Canada. Rapid responses are based on a limited literature search and are not comprehensive, systematic reviews. The intent is to provide a list of sources and a summary of the best evidence on the topic that CADTH could identify using all reasonable efforts within the time allowed. Rapid responses should be considered along with other types of information and health care considerations. The information included in this response is not intended to replace professional medical advice, nor should it be construed as a recommendation for or against the use of a particular health technology. Readers are also cautioned that a lack of good quality evidence does not necessarily mean a lack of effectiveness particularly in the case of new and emerging health technologies, for which little information can be found, but which may in future prove to be effective. While CADTH has taken care in the preparation of the report to ensure that its contents are accurate, complete and up to date, CADTH does not make any guarantee to that effect. CADTH is not liable for any loss or damages resulting from use of the information in the report.

Copyright: This report contains CADTH copyright material. It may be copied and used for non-commercial purposes, provided that attribution is given to CADTH.

Links: This report may contain links to other information available on the websites of third parties on the Internet. CADTH does not have control over the content of such sites. Use of third party sites is governed by the owners' own terms and conditions.

The aim of this review is to evaluate the comparative clinical effectiveness of these new machines or any other NPWT device at different pressures for the treatment of patients with diabetic foot or pressure ulcers.

RESEARCH QUESTIONS

1. What is the comparative clinical effectiveness of negative pressure wound therapy at -75 mmHg, -80 mmHg, and -125 mmHg for patients with diabetic foot ulcers and pressure ulcers?
2. What is the comparative clinical effectiveness of continuous versus intermittent negative pressure wound therapy at -75 mmHg, -80 mmHg, or -125 mmHg for patients with diabetic foot ulcers and pressure ulcers?

KEY MESSAGE

Due to lack of evidence, the comparative clinical effectiveness of negative pressure wound therapy at different pressures or comparative clinical effectiveness of continuous versus intermittent negative pressure wound therapy for patients with diabetic foot ulcers and pressure ulcers remains to be established.

METHODS

Literature Search Strategy

A limited literature search was conducted on key resources including PubMed, The Cochrane Library (2012, Issue 5), University of York Centre for Reviews and Dissemination (CRD) databases, Canadian and abbreviated lists of major international health technology agencies, as well as a focused Internet search. No filters were applied to limit the retrieval by study type. Where possible, retrieval was limited to the human population. The search was also limited to English language documents published between January 1, 2007 and May 24, 2012.

Selection Criteria and Methods

One reviewer screened the titles and abstracts of the retrieved publications, and evaluated the full-text publications for the final article selection, according to the selection criteria presented in Table 1.

Table 1: Selection Criteria

Population	Patients with diabetic foot ulcers or pressure ulcers
Intervention	Negative Pressure Wound Therapy
Comparator	Q1: Different pressures of NPWT (e.g., -75 mmHg, -80 mmHg, -125 mmHg) Q2: Continuous versus Intermittent NPWT
Outcomes	Wound healing
Study Designs	Health technology assessments, systematic reviews and meta-analysis, randomized controlled trials (RCTs) and non-randomized studies

Exclusion Criteria

Studies were excluded if they did not meet the selection criteria.

SUMMARY OF EVIDENCE

Quantity of Research Available

The literature search yielded 221 citations. Nine additional citations were identified in the grey literature. Upon screening titles and abstracts, 208 citations were excluded, and 22 potentially relevant articles were retrieved for full-text review. Of the 22 potentially relevant reports,^{1,3,6,12,13,21,27-42} no studies met the inclusion criteria. That is, no single study compared different pressures of NPWT (such as -75 mmHg, -80 mmHg, -125 mmHg, or other pressures) or compared continuous versus intermittent NPWT in patients with diabetic foot ulcers or pressure ulcers. The main reason for exclusion was lack of appropriate comparator (n = 15 studies). These studies compared NPWT with standard of care, rather than with another type of NPWT device or the same NPWT device operating at different pressures. The study selection process is outlined in a Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flowchart (Appendix 1).

CONCLUSIONS AND IMPLICATIONS FOR DECISION OR POLICY MAKING

No evidence was identified on the comparative clinical effectiveness of negative pressure wound therapy at -75 mmHg, -80 mmHg, and -125 mmHg, or other pressures, for the treatment of diabetic foot ulcers or pressure ulcers. Furthermore, no evidence on comparative clinical effectiveness of continuous versus intermittent negative pressure wound therapy in the management of these patients was identified. No trials were identified comparing different devices for the application of negative pressure wound therapy, including Smith & Nephew PICO or KCI V.A.C. Via devices. Well-designed clinical trials are needed for assessing the comparative clinical effectiveness of different pressures for NPWT for the treatment of diabetic foot ulcer and pressure ulcers.

PREPARED BY:

Canadian Agency for Drugs and Technologies in Health

Tel: 1-866-898-8439

www.cadth.ca

REFERENCES

1. Karatepe O, Eken I, Acet E, Unal O, Mert M, Koc B, et al. Vacuum assisted closure improves the quality of life in patients with diabetic foot. *Acta Chir Belg*. 2011 Sep;111(5):298-302.
2. Margolis D, Malay DS, Hoffstad OJ, Leonard CE, MaCurdy T, Lopez de Nava K, et al. Prevalence of diabetes, diabetic foot ulcer, and lower extremity amputation among Medicare beneficiaries, 2006 to 2008. *Diabetic foot ulcers. Data points #1* [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2011 Feb. [cited 2012 Jun 11]. (AHRQ publication no. 10(11)-EHC009-EF). Available from: http://effectivehealthcare.ahrq.gov/ehc/products/225/508/Data-Points_1_Diabetic-Foot-Ulcers_Report_02-2011.pdf Prepared by the University of Pennsylvania DEClIDE Center, under contract no. HHS290200500411
3. Centre for Clinical Practice at NICE. *Diabetic foot problems: inpatient management of diabetic foot problems* [Internet]. London (UK): National Institute for Health and Clinical Excellence (NICE); 2011 Mar. [cited 2012 Jun 11; updated 2012 Jan]. (NICE clinical guideline 119). Available from: <http://www.nice.org.uk/nicemedia/live/13416/53556/53556.pdf>
4. Gottrup F, Apelqvist J. Present and new techniques and devices in the treatment of DFU: a critical review of evidence. *Diabetes Metab Res Rev*. 2012 Feb;28 Suppl 1:64-71.
5. Reddy M. Pressure ulcers. *Clin Evid (Online)*. 2011 Apr 28;2011. pii:1901.
6. Effective Health Care Program. Evidence-based Practice Center systematic review protocol pressure ulcer treatment strategies: a comparative effectiveness review [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2008 Nov 8. [cited 2012 Jun 11]. Available from: http://effectivehealthcare.ahrq.gov/ehc/products/308/838/Pressure-Ulcer-Treatment_%20Protocol_20111108.pdf
7. Bergstrom N, Bennett MA, Carlson CE. Pressure ulcer treatment. Clinical practice guideline no. 15. Rockville (MD): Agency for Health Care Policy and Research (AHRQ); 1994. (AHCPR publication 95-0653).
8. Graves N, Birrell F, Whitby M. Effect of pressure ulcers on length of hospital stay. *Infect Control Hosp Epidemiol*. 2005 Mar;26(3):293-7.
9. Lyder CH. Pressure ulcer prevention and management. *JAMA*. 2003 Jan 8;289(2):223-6.
10. Nain PS, Uppal SK, Garg R, Bajaj K, Garg S. Role of negative pressure wound therapy in healing of diabetic foot ulcers. *J Surg Tech Case Rep* [Internet]. 2011 Jan [cited 2012 Jun 5];3(1):17-22. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3192517>
11. Nather A. Role of negative pressure wound therapy in healing of diabetic foot ulcers. *J Surg Tech Case Rep*. 2011 Jan;3(1):10-1.
12. de Laat EH, van den Boogaard MH, Spauwen PH, van Kuppevelt DH, van Goor H., Schoonhoven L. Faster wound healing with topical negative pressure therapy in difficult-to-

- heal wounds: a prospective randomized controlled trial. *Ann Plast Surg.* 2011 Dec;67(6):626-31.
13. Akbari A, Moodi H, Ghiasi F, Sagheb HM, Rashidi H. Effects of vacuum-compression therapy on healing of diabetic foot ulcers: randomized controlled trial. *J Rehabil Res Dev* [Internet]. 2007 [cited 2012 Jun 5];44(5):631-6. Available from: <http://www.rehab.research.va.gov/jour/07/44/5/pdf/akbari.pdf>
 14. Suissa D, Danino A, Nikolis A. Negative-pressure therapy versus standard wound care: a meta-analysis of randomized trials. *Plast Reconstr Surg.* 2011 Nov;128(5):498e-503e.
 15. Mandal A. Role of topical negative pressure in pressure ulcer management. *J Wound Care.* 2007 Jan;16(1):33-5.
 16. Xie X, McGregor M, Dendukuri N. The clinical effectiveness of negative pressure wound therapy: a systematic review. *J Wound Care.* 2010 Nov;19(11):490-5.
 17. Glass GE, Nanchahal J. The methodology of negative pressure wound therapy: separating fact from fiction. *J Plast Reconstr Aesthet Surg.* 2012 Jan 9. Epub ahead of print.
 18. Armstrong DG, Marston WA, Reyzelman AM, Kirsner RS. Comparative effectiveness of mechanically and electrically powered negative pressure wound therapy devices: a multicenter randomized controlled trial. *Wound Repair Regen.* 2012 May;20(3):332-41.
 19. Thompson G. An overview of negative pressure wound therapy (NPWT). *Br J Community Nurs.* 2008 Jun;13(6):S23-S30.
 20. Fogg E. Best treatment of nonhealing and problematic wounds. *JAAPA* [Internet]. 2009 Aug [cited 2012 Jun 14];22(8):46, 48. Available from: http://media.haymarketmedia.com/documents/10/whatsnew0809_2281.pdf
 21. Campbell PE, Smith GS, Smith JM. Retrospective clinical evaluation of gauze-based negative pressure wound therapy. *Int Wound J.* 2008 Jun;5(2):280-6.
 22. Topical negative pressure for chronic wounds? *Drug Ther Bull.* 2007 Aug;45(8):57-61.
 23. Borgquist O, Ingemansson R, Malmso M. The influence of low and high pressure levels during negative-pressure wound therapy on wound contraction and fluid evacuation. *Plast Reconstr Surg.* 2011 Feb;127(2):551-9.
 24. Borgquist O, Ingemansson R, Malmso M. Wound edge microvascular blood flow during negative-pressure wound therapy: examining the effects of pressures from -10 to -175 mmHg. *Plast Reconstr Surg.* 2010 Feb;125(2):502-9.
 25. Clinical data indicates PICO™ NPWT System reduces healing time and increases patient comfort [Internet]. [London]: Smith & Nephew; 2012 May 23. [cited 2012 Jun 11]. Available from: http://ewma2012.org/fileadmin/user_upload/EWMA/pdf/EWMA_2012/Press_Release_SmithNephew_Pico_01.pdf

26. V.A.C.Via™ therapy system: monograph [Internet]. [San Antonio, TX]: KCI; 2011. [cited 2012 Jun 11]. Available from:
<http://www.kci1.com/cs/Satellite?blobcol=urldata&blobheadname1=Content-type&blobheadname2=Content-disposition&blobheadname3=MDT-Type&blobheadvalue1=application%2Fpdf&blobheadvalue2=inline%3B+filename%3D17%252F213%252FV.A.C.Via%25E2%2584%25A2%2BTherapy%2BSystem%2BMonogr aph.pdf&blobheadvalue3=abinary%3B+charset%3DUTF-8&blobkey=id&blobtable=MungoBlobs&blobwhere=1226650908227&ssbinary=true>
27. Blume PA, Walters J, Payne W, Ayala J, Lantis J. Comparison of negative pressure wound therapy using vacuum-assisted closure with advanced moist wound therapy in the treatment of diabetic foot ulcers: a multicenter randomized controlled trial. *Diabetes Care*. 2008 Apr;31(4):631-6.
28. Masden D, Goldstein J, Endara M, Xu K, Steinberg J, Attinger C. Negative pressure wound therapy for at-risk surgical closures in patients with multiple comorbidities: a prospective randomized controlled study. *Ann Surg*. 2012 Jun;255(6):1043-7.
29. Khanbhai M, Fosah R, Oddy MJ, Richards T. Disposable NPWT device to facilitate early patient discharge following complex DFU. *J Wound Care*. 2012 Apr;21(4):180, 182.
30. Rahmanian-Schwarz A, Willkomm LM, Gonser P, Hirt B, Schaller HE. A novel option in negative pressure wound therapy (NPWT) for chronic and acute wound care. *Burns*. 2012 Jun;38(4):573-7.
31. Ulusal AE, Sahin MS, Ulusal B, Cakmak G, Tuncay C. Negative pressure wound therapy in patients with diabetic foot. *Acta Orthop Traumatol Turc*. 2011;45(4):254-60.
32. Bondokji S, Rangaswamy M, Reuter C, Farajalla Y, Mole T, Cockwill J, et al. Clinical efficacy of a new variant of a foam-based NWPT system. *J Wound Care*. 2011 Feb;20(2):62, 64-5-67.
33. Stansby G, Wealleans V, Wilson L, Morrow D, Gooday C, Dhatariya K. Clinical experience of a new NPWT system in diabetic foot ulcers and post-amputation wounds. *J Wound Care*. 2010 Nov;19(11):496, 498-9-502.
34. Witkowski W, Jawien A, Witkiewicz W, Zon B. Initial multi-centre observations upon the effect of a new topical negative pressure device upon patient and clinician experience and the treatment of wounds. *Int Wound J*. 2009 Apr;6(2):167-74.
35. Fife CE, Walker D, Thomson B, Otto G. The safety of negative pressure wound therapy using vacuum-assisted closure in diabetic foot ulcers treated in the outpatient setting. *Int Wound J*. 2008 Jun;5 Suppl 2:17-22.
36. van den Boogaard M, de Laat E, Spauwen P, Schoonhoven L. The effectiveness of topical negative pressure in the treatment of pressure ulcers: a literature review. *Eur J Plast Surg* [Internet]. 2008 [cited 2012 Jun 11];31(1):1-7. Available from:
<http://www.springerlink.com/content/w611780x8v74j147/fulltext.html>
37. Xie X, McGregor M. Negative pressure wound therapy (NPWT) (update of report 19) [Internet]. Report no. 48. Montreal: Technology Assessment Unit (TAU) of the McGill

- University Health Centre (MUHC); 2010 Jun 25. [cited 2012 Jun 11]. Available from: https://secureweb.mcgill.ca/tau/sites/mcgill.ca/tau/files/NPWT_FINAL.pdf
38. Medical Advisory Secretariat. Negative pressure wound therapy: an evidence update. Ont Health Technol Assess Ser [Internet]. 2010 Dec [cited 2012 Jun 11];10(22):1-28. Available from: http://www.health.gov.on.ca/english/providers/program/mas/tech/reviews/pdf/update_NPWT_20101214.pdf
39. ECRI Institute, Sullivan N, Snyder DL, Tipton K, Uhl S, Schoelles KM. Negative pressure wound therapy devices: technology assessment report [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2009 Nov 12. [cited 2012 Jun 11]. Available from: <http://www.ahrq.gov/clinic/ta/negpresswtd/negpresswtd.pdf>
40. Ubbink DT, Westerbos SJ, Evans D, Land L, Vermeulen H. Topical negative pressure for treating chronic wounds. Cochrane Database Syst Rev. 2008 Jul 16;(3):CD001898.
41. Swedish Council on Technology Assessment in Health Care (SBU), The Regional HTA Centre of Region Västra Götaland. Vacuum-assisted wound closure after surgical interventions [Internet]. Stockholm: SBU; 2011 [cited 2012 Jun 11]. (SBU Alert report no. 2011-09). Available from: http://www.sbu.se/upload/Publikationer/Content1/1/Vacuum_Assisted_Wound_Closure_Therapy.pdf
42. Fong KD, Hu D, Eichstadt SL, Gorell E, Munoz CA, Lorenz HP, et al. Initial clinical experience using a novel ultraportable negative pressure wound therapy device. Wounds [Internet]. 2010 [cited 2012 Jun 11];22(9):230-6. Available from: <http://www.woundsresearch.com/files/wounds/pdfs/fong.pdf>

APPENDIX 1: Selection of Included Studies

